

1. To Improve accuracy of patient identification use at least two patient identifiers. List two acceptable identifiers.
  
2. Which clinical situations would the nurse always compare the arm band with the request slip? Check all that apply:
  - Providing care, treatment, or services
  - Administering medications or blood products
  - Taking blood samples and other specimens for clinical testing
  - Starting an IV
  
3. The unit nurse has just received a telephone order from a primary care provider. To improve the effectiveness of communication among caregivers the nurse will follow a sequence of critical steps when **taking an order** to ensure patient safety. Please list these steps :
  
4. To reduce the risk of health care associated infections all care providers perform hand hygiene (for minimum 15 seconds) prior to and after patient care. Soap and water must be used when caring for patients with \_\_\_\_\_ to kill the spores. In addition, the nurse should place patients with infectious conditions in \_\_\_\_\_.
  
5. To reduce the risk of patient harm resulting from falls identify when a fall risk assessment should be initiated, and where is it documented?
  
6. List the 'Universal' Fall Precautions the nurse would implement for **all** patients:
  
  
7. To reduce errors in giving high risk medication, **circle the items/meds** that must be double-checked by another nurse or pharmacist:  
**Chemo, Insulin, Bolus anticoagulants, Blood products, PCA meds/pumps, Pediatric meds**